

## Edward-Elmhurst Cancer Centers

120 Spalding Drive; Suite 111; Naperville, IL 60540 Phone: 630/646-2273 Fax: 630/548-6617

24600 W. 127<sup>th</sup> Street; Plainfield, IL 60585 Phone: 630/646-2273 Fax: 630/548-6617 177 E. Brush Hill Road; Elmhurst, IL 60126 Phone: 630/646-2273 Fax: 331/221-3887

	Actemra	(TOCILIZUMAB) Standing Order					
Patient Name:		DOB:					
***Please include current history and physical and any recent labs/tests, if applicable.*							
*PLEASE A	ATTACH COPY	OF INSURANCE CARD WITH THIS ORDER*					
Pre-Authorization Reference #:		ysician Office is Responsible to Obtain Authorization/Referral)					
Contact Name an Number of Insura							
If you have any qu billing department.	• • • • • • • • • • • • • • • • • • • •	e-authorizations, please contact (630) 527-3788 and ask for the					
Allergies:							
Diagnosis (ICD 10	Required):						
Weight:	Lbs:	Kg:					
inadequate respons	se to 1 or more TNF a	ult patients with moderately to severely active RA who had an antagonist therapies. Recommended starting dose is 4mg/kg ery 4 weeks based on clinical response.					
Pre-administratio	Current TB skin tes ANC, Platelet cour Actemra if ANC < 2 liver enzymes > 1.5 Pre-Infusion asses	st or chest x-ray  nt, Liver enzymes (ALT, AST) (Not recommended to start 2000/mm3, platelets < 100,000/mm3 and 5 times upper limit of normal) ssment to include BP, temp, pulse per unit protocol ssment to include BP, temp, pulse per unit protocol					
	ne following labs (ple CBC w/platelets an Liver Enzymes Other: equency (please ch	nd Auto Differential					
	Every 4 weeks dor	ne One week prior to infusion ne One week prior to infusion					

	☐ Each Infusion ☐ Other:										
•											
2.	<ol> <li>Please draw the following labs (please checking in the following labs).</li> <li>Lipid Panel</li> <li>Other:</li> </ol>										
	Desired Frequency (please check):										
	One week prior to first infusion, then 4 weeks after treatment initiation, then ever										
	6 months	sian than Quarks often treatme	ant initiation, then avenue								
	One week prior to first infus	sion, then 8 weeks after treatm	ent initiation, then every								
	<b>—</b>	tiation, then every 6 months									
	Other:										
Ad	Administration:										
1.		mg of Actemra ever	y 4 weeks.								
	Maximum dose is 800mg										
2.	2. Infuse 100ml Actemra solution over 6	60 minutes. Do not administer	as bolus or push.								
3.	B. Do not administer Actemra during an	active infection, including loca	lized infections.								
4.	Interrupt Actemra:	Interrupt Actemra:									
	If a serious infection develo	ops, including localized infectio	ns								
	ANC 500-1000	000									
	☐ Platelet count 50,000-100,0	บบบ LN after modification of DMAR	D dose (4ma/ka dose)								
	Liver Enzymes > 3 to 5X U		b dood ( img/kg dood)								
5.	5. Discontinue Actemra:										
5.	ANC < 500										
	☐ Platelets < 50,000										
	☐ Liver enzymes > 5X ULN										
6	6. Reduce Actemra dose from 8mg/kg to 4mg/	/ka·									
Ο.		LN after modification of DMAR	D dose								
_											
1.	<ol><li>Do not administer Actemra concomitantly in the same line with other drugs. Avoid use of Ac in combination with other biologic DMARDS.</li></ol>										
	in combination with other blologic binArtbe	<b>.</b>									
8.	3. Infusion reactions may include:										
Ο.	•	lives/pruritis	*Hypertension								
	*Shortness of breath *Ba	ackache	*Dizziness *Anaphylaxis								
	*Nausea/vomiting *C	hest tightness									
9.	9. In the event of a hypersensitivity reaction	n during the infusion of this	medication, we will								

- 9. In the event of a hypersensitivity reaction during the infusion of this medication, we will implement the reaction protocol. A designated nurse practitioner will evaluate your patient and your office will receive notification of the event.
- 10. In the event that your patient has a central line, it will be used per the Cancer Center protocol, unless otherwise directed.

Revision/Review Date: 12/17/24											
Physician Name (Please Print)	O1	Office Phone		Fax Number							
Ordering Physician NPI:		_ Swedish Hospital N	IPI:	1831	15125	57					
Physician Signature:		Date:									
12. Patient may be discharged post-infusion if stable. No monitoring time required.											
<ul> <li>Methylprednisolone 100mg IV Push</li> </ul>	IV Push		Yes		No						
Acetaminophen	mg	PO	Yes		No						
<ul><li>11. Premedicate with:</li><li>Benadryl</li></ul>	mg	PO or IV (circle one)	Yes		No						